

| REPORT DOCUMENTATION PAGE | | | | Form Approved OMB No. 0704-0188 | |
|---|-----------------------------|--|--------------------------------------|---|---|
| The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION. | | | | | |
| 1. REPORT DATE (DD-MM-YYYY) 15-02-2012 | | 2. REPORT TYPE Conference Proceedings | | 3. DATES COVERED (From - To) | |
| 4. TITLE AND SUBTITLE Evaluation of Wind Field Predictions by Atmospheric Models Over the Marmara Sea | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER 0602435N | |
| 6. AUTHOR(S) Mustafa Kemal Cambazoglu, Cheryl Ann Blain, M. Kadioglu | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER 73-6263-09-5 | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7320-09-0091 | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research One Liberty Center 875 North Randolph Street, Suite 1425 Arlington, VA 22203-1995 | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) ONR | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited. | | | | | |
| 13. SUPPLEMENTARY NOTES <div style="text-align: center; font-size: 1.5em; margin-top: 20px;">20120217341</div> | | | | | |
| 14. ABSTRACT Data collected from meteorological stations in the Turkish Straits System around the Marmara Sea are used to assess the performance of atmospheric models in predicting the winds. The Coupled Ocean/Atmosphere Mesoscale Prediction System, COAMPS is applied using different spatial resolutions ranging between 1 km and 27 km to investigate the effect of model spatial grid resolution on accuracy of computed wind field. The influence of the ocean dynamics on the atmospheric winds is also investigated by comparing wind field predictions from a fully coupled COAMPS with those from the uncoupled (stand-alone atmospheric) COAMPS. Following the examination of the wind products, the importance of using high resolution wind forcing for ocean circulation predictions is evaluated. | | | | | |
| 15. SUBJECT TERMS Marmara Sea, air-sea interactions, wind/foat, circulation, atmospheric input | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT UU | 18. NUMBER OF PAGES 1 | 19a. NAME OF RESPONSIBLE PERSON Cheryl Ann Blain |
| a. REPORT Unclassified | b. ABSTRACT Unclassified | c. THIS PAGE Unclassified | | | 19b. TELEPHONE NUMBER (Include area code) 228-688-5450 |

EVALUATION OF WIND FIELD PREDICTIONS BY ATMOSPHERIC MODELS OVER THE MARMARA SEA

Mustafa Kemal Cambazoglu^{1*}, Cheryl Ann Blain² and Mikdat Kadioglu³

¹ Department of Marine Science, The University of Southern Mississippi, Stennis Space Center, MS, USA - kemal.cambazoglu.ctr.tu@nrlssc.navy.mil

² Oceanography Division, Naval Research Laboratory, Stennis Space Center, MS, USA

³ Faculty of Aeronautics and Astronautics, Istanbul Technical University, Istanbul, Turkey

Abstract

Data collected from meteorological stations in the Turkish Straits System around the Marmara Sea are used to assess the performance of atmospheric models in predicting the winds. The Coupled Ocean/Atmosphere Mesoscale Prediction System, COAMPS is applied using different spatial resolutions ranging between 1 km and 27 km to investigate the effect of model spatial grid resolution on accuracy of the computed wind field. The influence of the ocean dynamics on the atmospheric winds is also investigated by comparing wind field predictions from a fully coupled COAMPS with those from the uncoupled (stand-alone atmospheric) COAMPS. Following the examination of the wind products, the importance of using high resolution wind forcing for ocean circulation predictions is evaluated.

Keywords: *Marmara Sea, Air-Sea Interactions, Wind/Font, Circulation, Atmospheric Input*

The Turkish Straits System (TSS) contains the Dardanelles and the Bosphorus straits that connect the Marmara Sea to the Aegean and the Black Seas, respectively. Fresh, light Black Sea water flows at the surface towards the Aegean Sea while saline, dense Mediterranean water flows deep in the opposite direction resulting in a two-layer stratified flow in both straits. Winds over the region play a major role in generating various features within the TSS system, i.e. blocking of the deep (surface) flow within the straits by Northerly (Southerly) winds; the spread of the Dardanelles outflow plume over the Aegean Sea; and the circulation pattern within the Marmara Sea among the others. Since the winds are the dominant external forcing applied to the ocean models, accurate predictions of winds by atmospheric models are crucial for accurate predictions of the flow dynamics.

Hourly wind direction and speed data collected from 13 meteorological stations located around the Marmara Sea between May 2008 and June 2009 are used to assess the performance of atmospheric models in predicting winds. The measurements recorded every ten minutes by two Meteo-buoys on the Marmara and Black Seas are also included in the evaluation. The atmospheric models under consideration are the Coupled Ocean/Atmosphere Mesoscale Prediction System, COAMPS [1], and the numerical weather prediction model, COSMO-ME, based on the non-hydrostatic Lokal-Model(LM) [2]. COSMO is the Consortium for Small scale Modeling and COSMO-ME is a particular application running on the Southern Europe/Mediterranean Sea domain managed by USAM (Meteorological Service of the Italian AirForce). It has a spatial resolution of 7-km while COAMPS has a coarse operational grid resolution of 27-km. Local configurations of the COAMPS model for grids of 9-km, 3-km and 1-km are also considered.




Fig. 1. Wind speed variation (36-hour averaged) over September and October 2008 at Canakkale and Bandirma meteorological station locations: Measurements (solid), 7-km COSMO model results (dash) and 27-km COAMPS model results (dot).




Fig. 2. Frequency and coherence spectra of wind direction measurements vs. the COAMPS (27-km) and COSMO (7-km) model results for Canakkale and Bandirma meteorological stations.

Both model inter-comparisons (COAMPS vs. COSMO) and model-data comparisons are conducted. Model-data comparisons consider both the raw and filtered time series of wind speed and direction. A more detailed statistical analysis is done by examining the frequency spectra of the modeled and measured wind fields and the coherence between them. The coarser COAMPS model is able to capture the low-frequency (36-hour averaged) wind speed variability as shown in Fig. 1. COAMPS-COSMO comparisons indicate similar wind field predictions (shown in Fig.1) with the exception of wind direction during higher frequency wind events such as the semi-diurnal cycle or the land/sea breeze events as shown in Fig. 2.

Acknowledgements

This work is supported by the Office of Naval Research under the 6.2 NRL program "Development of a Multi-Scale Coupled Ocean Model System: Application to the Turkish Straits". The authors appreciate Dr. E. Jarosz of NRL for the software that facilitated model-data coherence analysis as well as Drs. J. Dykes and T. Smith of NRL for providing the model results of locally configured COAMPS simulations. Further thanks are extended also to Dr. J. Chiggiato of NURC for providing the COSMO-ME data for the evaluations. This paper is contribution number NRL/PP/7320--09-0091

References

- 1 - Hodur, R. M., 1997. The Naval Research Laboratory's coupled ocean/atmosphere mesoscale prediction system (COAMPS). *Mon. Weather Rev.*, 125(7): 1414-1430
- 2 - Steppeler J., Doms G., Schättler U., Bitzer H.W., Gassmann A., Damrath U. and Gregoric G., 2003. Meso-gamma scale forecasts using the nonhydrostatic model LM. *Meteorol. Atmos. Phys.*, 82: 75-96.